

Locating possible micro-osteoperforation sites and alveolar bone thickness by CBCT in young adults

By Afrah Khazal Al Hamdany

Locating possible micro-osteoperforation sites and alveolar bone thickness by CBCT in young adults

Afrah Khazal Al Hamdany, Tara Ali Rasheed, Nada Nashee Alhafidh

¹³ Department of Pedodontics, Orthodontics, and Preventive Dentistry, College of Dentistry, Mosul University, Iraq

Tara Ali Rasheed
<https://orcid.org/0000-0001-6730-5114>

Nada Nashee Alhafidh
<https://orcid.org/0000-0002-1853-5826>

²⁵ e-mail: afrah2@uomosul.edu.iq
<https://orcid.org/0000-0002-8316-0215>

ABSTRACT

Objectives, Micro-osteoperforation (MOP) was introduced as a minimum invasive surgical intervention employed ³² to accelerate orthodontic tooth movement. This study aimed to determine proposed MOP sites with the available ³⁶ alveolar bone thickness (ABT) needed in the maxilla (Max.) and the mandible (Mand.) at both genders ¹¹ using cone-beam computed tomography (CBCT).

Methods. Pre-operative CBCT scans of 48 skeletal Class I (18-30yrs old) patients (22 males, 26 females) were analyzed. NewTom's NNT software was used to measure ABT. Measurements were done in 13 bilateral Max. and Mand. inter-radicular sites at 3 levels corresponding to the proposed

MOP sites. ²⁸ Statistical Package for Social Sciences, version 21 was used. ³⁴ Descriptive statistic values and Independent samples t-test were used to test the gender and Max.-Mand. ABT difference, $P \leq 0.05$.

Results. There was a general tendency toward increasing ABT gradually from the inter-incisal region distally toward the molar and from near cement-enamel junction toward the apical region in Max. arch in both genders. In Mand., the same trend was observed as in Max., especially for the males. No statistically significant gender ABT difference found at both dental arches except those of females who had significantly higher Max. ABT at specific locations. ²⁴ No statistically significant difference was found in the comparison between both dental arches at $P \leq 0.05$ except for certain locations where the Max. had significantly larger values than Mand.

Conclusions. detailed information for ABT concerning MOP sites was obtained at different locations of both dental arches for both genders. Abundant ABT was found apically at Max. posterior region.

Key words: ¹² Cone-beam computed tomography, Micro-osteoperforations, Orthodontics, Tooth movement.

INTRODUCTION

The long duration of orthodontic treatment is one of the chief problems facing both the patient and the orthodontist [1], which can raise the severity and the risk of the originated adverse effects from orthodontic therapy [2]. Since the start of orthodontic treatment, many interventions have been made to curtail the total time of treatment, employing diverse appliances and approaches with variable success rates. The non-surgical approaches include electromagnetic fields [3], custom-made brackets and wires [4], self-ligating brackets [5], medications [6],

low-level laser and photodynamic therapy [7], low-intensity high-rate vibrations [8] and cell mediators injection [9].

Some evidences; obtained by ⁸ meta-analysis and randomized controlled trials; indicated that the surgical approaches may be more effective in terms of accelerating the rate of tooth movement [10]. Invasive surgical approaches, ⁵ such as osteotomies [11], corticotomies [12] associated with or without a bone graft, and less invasive approaches as piezocisions [13], piezopuncture and micro-osteoperforation (MOP) [14] have been conducted for stimulating the bone natural mechanisms [15], increasing the inflammatory mediators; triggered by orthodontic forces [16]; that temporarily upgrade the bone metabolism and resorption and could possibly affect the tooth movement rate [17].

¹ Among the recent modalities introduced to accelerate orthodontic tooth movement is MOP, in other words, bone puncturing using mini-screw or propel device ³⁷ to induce micro-trauma to the alveolar bone initiating a regional acceleratory phenomenon [18]. MOP; ¹ has been introduced to meet the growing need for accelerating orthodontic treatment, especially by adult patients; is an approach based on principles of sound bone biology. It is a minimally invasive, safe approach ¹ that can be employed in conjunction with any orthodontic appliances and can be performed by the orthodontist, ¹⁸ not only to accelerate orthodontic tooth movement with no or less pain or discomfort [14,19,20], but in many other orthodontic situations, obviously for changing the tooth movement type or creating differential anchorage [21]. According to Aboalnaga ¹⁷ *et al.* [22], Alqadasi *et al.* [23], Fattoria *et al.* [24] and Mittal *et al.* [25], MOPs were not able to accelerate the tooth movement rate; nevertheless, ³⁵ they seemed to assist root movement. Shahabee *et al.* [26] in their ⁵ systematic review and meta-analysis evaluated the effects of MOP on the

tooth movement rate emphasizing that MOP can be suggested after weighing the benefits and drawbacks of this approach.

The determination of alveolar bone thickness (ABT) is a critical step in predicting treatment outcomes related to MOP procedures. No previous study in the literature was found with the detailed ABT at different locations in the maxilla (Max.) and mandible (Mand.) for both genders, specially concerning MOP sites. Thus, the objective of the current study was to determine Max. and Mand. ABT; especially concerning the determination of proposed MOP sites using a high-resolution CBCT system in the young adult population; at 3, 5 and 7 mm level from cemento-enamel junction (CEJ) and to reveal the gender and Max.-Mand. ABT differences. The tested hypothesis was that there were no gender or Max.-Mand. differences in ABT.

MATERIALS AND METHODS

Study protocol and registration: The study protocol was registered and approved (at session No. UoM.Dent/H.68/22) by the Research Ethics Committee of the College of Dentistry/ University of Mosul/Iraq and (at session No. 193/23; meeting No.48) by the Research Ethics Committee of the College of Dentistry/ University of Sulaimani /Iraq.

Study design and sample characteristics

This was a retrospective observational study and the sample represented CBCT scans of orthodontic patients that were previously treated orthodontically. The selected CBCT scans were chosen according to the next eligibility criteria:

Inclusion Criteria

Eastern Mediterranean young adult Kurdish (18-30 years old) patients having Class I skeletal relationship, with upper and/or lower dental arch length discrepancy of up to 6mm. CBCT scans were presented including the nasal bone to the chin area. All CBCT scans had been gotten with maximum intercuspation.

Exclusion Criteria

Congenitally missing or extracted teeth except for third molars, incomplete erupted teeth or impacted teeth, supernumerary teeth, facial asymmetry, craniofacial deformity, horizontal or vertical alveolar bone loss due to periodontal disease seen in CBCT scans, restorations involving CEJ and orthodontic treatment history.

After applying selection criteria on 89 CBCT scans, 48 scans (22 for males and 26 for females) with mean age of 29.25 and 26.5 years, respectively, were involved in this study. The following formula was used for sample size calculation: $n = 2\sigma^2 / \Delta^2 (z\alpha + z\beta)^2$. The resulted number was adjusted, and the final sample size in each group = $n + (n \times 0.2)$. In this study, n is considered as the number of subjects, σ (standard deviation) = 1.22 mm (representing the standard deviation of Max.ABT according to Andrews et al. [27]), (Δ) precision = 0.2 unit, $z\alpha = 1.96$ for $\alpha = 0.05$ and $z\beta = 0.84$ for 80% power. According to this, for each gender group, the calculated sample size was approximately 11 subjects. And for more accurate and precise results, larger than this number was used in each group.

The scans were previously performed using Cone Beam Tomography NewTom machine (NewTom VGi scanner; QR s.r.l., Verona, Italy). NewTom CBCT machine was set at Full Scale Voltage

(FSV) 90 KV, 50.25 mAs, with 0.15 mm nominal slice thickness, field of view (FOV) 8*5 HRes, and exposure time 9.0s.

Measurements

NewTom’s NNT software was used to measure ABT. After the software was opened, CBCT image was downloaded. We chose the multi-planar option from the toolbar, and four views coordinated with each other appeared on a screen including: axial, panoramic, sagittal and 3D rendering images [Figure 1-A]. The axial and panoramic views were used to determine the inter-radicular site intended to be measured, and the sagittal view that was highly outlined with red color was used to measure ABT. For each CBCT image, bilateral measurements were done for ABT in 13 Max. inter-radicular sites from the mesial site of the right (R) second molar to the distal site of the left (L) first molar and 13 bilateral Mand. sites from the mesial site of the L second molar to the distal site of the R first molar corresponding to the proposed MOP sites. For each site, ABT was measured perpendicularly from the buccal cortical plate to the lingual/palatal cortical plate at 3, 5 and 7 mm levels from CEJ parallel to it [Figure 1-B]. All measurements were recorded in millimeters.




Figure 1: NewTom's NNT software for CBCT images. A: showed four views include: Axial, Panoramic, Sagittal and 3D rendering images, B: Bone thickness measured in sagittal view at 3,5 and 7 mm from cemento-enamel junction.

All the measurements were accomplished by one observer. Intra- and inter-class reliability correlations were calculated using 10 randomly selected CBCT images.

9 Statistical analyses

The measurements were analyzed using SPSS (Statistical Package for Social Sciences), version 21 (SPSS Inc., Chicago, IL) as follows: Data was firstly inspected for normal distribution using Shapiro-Wilk's normality test. The mean and standard deviation values were calculated. An independent samples t-test was used to test the gender and Max.-Mand. differences in ABT. The level of significance was $P \leq 0.05$.

RESULTS

For all measured parameters, the inter- and intra-class reliability were (>0.8) representing good inter- and intra-observer reliability. Shapiro-Wilk's Test revealed a normal distribution of data at $P \leq 0.05$. Thus parametric statistics were followed.

Descriptive statistics

Tables [1] and [2] revealed the descriptive statistics of Max.-Mand. ABT at different levels from CEJ for the males and females. The least value was for males at Mand. anterior region Mesial to R1 at 3mm level (5.09 ± 0.87), while the highest value was for Males at Max. posterior region distal to L6 at 7mm level (14.23 ± 1.57).

For males and females, the mean bilateral values of Max. ABT at the anterior, the premolar and at the molar regions were observed at Figure[2]. There is a tendency toward increasing in Max. ABT gradually from inter-incisal distally toward molar and from near CEJ toward apical regions, in both genders,

The mean bilateral values of Mand. ABT of males and females at the anterior, the premolar and at the molar regions were noticed at Figure [3]. Also, there is a tendency toward increasing Mand. ABT gradually from inter-incisal distally toward molar and from near CEJ toward apical regions, especially in the males.

Table 1: Descriptive statistics of maxillary alveolar bone thickness at different levels from cement-enamel junction for males and females with gender comparison

Site	Level (mm)	Males	Females	Sig.
		Mean (SD)	Mean (SD)	
Mesial to R7	3	12.63 (1.54)	13.99 (1.06)	S
	5	13.55 (0.92)	14.48 (0.95)	S
	7	13.87 (1.29)	14.70 (0.91)	NS
Mesial to R6	3	10.47 (1.27)	10.75 (1.06)	NS
	5	10.83 (1.15)	11.18 (0.70)	NS
	7	11.31 (1.29)	11.43 (0.86)	NS
Mesial to R5	3	8.75 (0.83)	8.93 (1.34)	NS
	5	9.21 (0.80)	9.59 (1.10)	NS
	7	9.18 (0.75)	9.90 (0.99)	NS
Mesial to R4	3	8.97 (1.37)	8.92 (1.26)	NS
	5	9.41 (1.23)	10.36 (1.18)	NS
	7	8.98 (1.51)	11.33 (2.05)	S
Mesial to R3	3	7.48 (1.11)	8.24 (1.40)	NS
	5	9.19 (1.94)	9.82 (2.24)	NS
	7	9.47 (2.40)	10.57 (2.61)	NS
Mesial to R2	3	8.90 (2.50)	8.76 (2.21)	NS
	5	9.82 (2.12)	10.61 (2.77)	NS
	7	10.12 (1.97)	11.16 (2.83)	NS
Mesial to	3	7.49 (1.68)	7.69 (1.03)	NS

R1	5	7.62 (2.34)	7.95 (1.09)	NS
	7	7.90 (2.68)	8.27 (1.40)	NS
Distal to L1	3	8.59 (1.93)	9.18 (1.97)	NS
	5	9.50 (1.82)	10.06 (2.28)	NS
	7	9.72 (1.99)	10.49 (1.77)	NS
Distal to L2	3	8.01 (2.25)	8.89 (1.87)	NS
	5	9.11 (2.08)	10.19 (2.07)	NS
	7	9.07 (2.32)	10.64 (2.61)	NS
Distal to L3	3	8.47 (1.06)	8.88 (1.20)	NS
	5	8.83 (0.91)	9.84 (1.19)	NS
	7	9.14 (1.57)	10.06 (1.73)	NS
Distal to L4	3	8.61 (1.14)	9.18 (0.98)	NS
	5	9.10 (1.39)	9.86 (1.11)	NS
	7	8.89 (1.56)	9.96 (1.18)	NS
Distal to L5	3	10.04 (0.48)	10.84 (0.71)	S
	5	10.77 (1.04)	11.17 (0.73)	NS
	7	10.79 (1.35)	10.98 (0.95)	NS
Distal to L6	3	13.32 (1.05)	13.70 (0.97)	NS
	5	13.91 (1.28)	14.41 (1.09)	NS
	7	14.23 (1.57)	14.17 (1.19)	NS

R=Right, L=Left, Max=Maxillary, Mand=Mandibular, **AB₉**= Alveolar Bone Thickness, SD=Standard Deviation, Males= 22, Females=26, Sig.= significant at $P < 0.05$.

Table 2: Descriptive statistics of mandibular alveolar bone thickness at different levels from cement-enamel junction for males and females with gender comparison

Site	Level (mm)	males	females	Sig.
		Mean (SD)	Mean (SD)	
Mesial to R7	3	10.49 (1.00)	11.86 (2.51)	NS
	5	11.84 (1.14)	13.73 (2.08)	S
	7	12.69 (1.95)	14.46 (1.62)	S
Mesial to R 6	3	8.91 (1.23)	8.67 (1.19)	NS
	5	9.81 (1.05)	10.09 (1.31)	NS
	7	10.41 (0.92)	10.90 (1.44)	NS
Mesial to R5	3	8.00 (1.38)	7.27 (0.81)	NS
	5	9.05 (1.66)	8.21 (1.29)	NS
	7	9.40 (2.03)	8.93 (1.59)	NS
Mesial to R4	3	7.90 (1.36)	7.16 (1.05)	NS
	5	9.31 (1.81)	8.16 (1.18)	NS
	7	9.62 (2.26)	8.55 (1.46)	NS
Mesial to R3	3	7.35 (1.82)	6.62 (3.44)	NS
	5	8.32 (1.60)	7.35 (1.18)	NS
	7	8.07 (1.81)	7.42 (1.46)	NS
Mesial to R2	3	6.90 (2.88)	6.03 (0.68)	NS
	5	6.75 (1.74)	6.77 (1.13)	NS
	7	6.54 (1.97)	6.55 (1.44)	NS
Mesial to R1	3	5.09 (0.87)	5.46 (0.78)	NS
	5	5.89 (1.40)	5.73 (1.01)	NS
	7	6.24 (1.36)	6.35 (1.09)	NS

Distal to L1	3	10.38 (1.43)	11.40 (1.61)	NS
	5	11.25 (1.56)	13.22 (1.78)	NS
	7	12.03 (1.77)	12.13 (2.63)	NS
Distal to L2	3	8.48 (1.16)	9.00 (0.96)	NS
	5	8.91 (1.50)	9.82 (1.02)	NS
	7	9.26 (1.69)	10.63 (1.05)	NS
Distal to L3	3	7.56 (1.48)	7.46 (0.92)	NS
	5	8.60 (1.76)	8.22 (1.29)	NS
	7	9.02 (1.51)	8.90 (1.46)	NS
Distal to L4	3	8.49 (1.14)	7.31 (1.63)	NS
	5	9.27 (1.48)	8.43 (1.62)	NS
	7	9.24 (1.64)	8.74 (1.58)	NS
Distal to L5	3	6.92 (0.94)	6.84 (1.31)	NS
	5	7.93 (1.48)	7.59 (1.38)	NS
	7	7.52 (1.67)	7.56 (1.57)	S
Distal to L6	3	6.47 (0.90)	6.26 (0.79)	NS
	5	7.10 (0.65)	6.86 (0.67)	S
	7	6.93 (1.14)	6.87 (0.74)	NS

R=Right, L=Left, Max=Maxillary, Mand=Mandibular, **AB6**= Alveolar Bone Thickness, SD=Standard Deviation, Males= 22, Females=26, Sig.= significant at $P \leq 0.05$, S= significant , NS= non significant.

Gender comparison

Tables [1] and [2] displayed significant statistical difference in ABT, in comparison between both gender, in the following sites: Max. Mesial to R7 at level of 3 and 5mm, Max. Mesial to R4 at level 7mm, Max. Distal to L5 at 3mm level, Mand. Mesial to R7 at level of 5 and 7mm and Mand. Distal to L5 at level 7mm in which the females have the largest value and Mand. Distal to L6 at a level of 5mm in which the males have the largest value. The rest sites were not statistically significant.




Figure 2: Bilateral maxillary mean alveolar bone thickness at different levels for males(A) and females(B).




Figure 3: Bilateral mandibular mean alveolar bone thickness at different levels for males(A) and females(B).

Maxillary-mandibular comparison

For males, Max.-Mand. comparison of ABT [Table 3] ³⁰ showed that there was significant difference in the following sites in which Max. was larger than Mand.: R and L Mesial to 7 and Mesial to 6 at three levels (except L Mesial to 7 and 6 at 7mm level); R Mesial to 5 at 3 and 5 levels; R Mesial to 3 at 7mm level; R and L Mesial to 2 (except at 3mm level at L side) and R and L Mesial to 1 at 3 and 5 mm levels. The rest sites were not statistically significant.

Table 3: Maxillary- mandibular comparison of alveolar bone thickness at different levels from cement-enamel junction for males

Variable	Level (mm)	R			L		
		Max. Mean (SD)	Mand. Mean (SD)	Sig.	Max. Mean (SD)	Mand. Mean (SD)	Sig.
Mesial to 7	3	12.63 (1.54)	10.38 (1.43)	S	13.32 (1.05)	10.49 (1.00)	S
	5	13.55 (0.92)	11.25 (1.56)	S	13.91 (1.28)	11.84 (1.14)	S
	7	13.87 (1.29)	12.03 (1.77)	S	14.23 (1.57)	12.69 (1.95)	NS
Mesial to 6	3	10.47 (1.27)	8.48 (1.16)	S	10.04(0.48)	8.91(1.23)	S
	5	10.83 (1.15)	8.91 (1.50)	S	10.77 (1.04)	9.81 (1.05)	S
	7	11.31 (1.29)	9.26 (1.69)	S	10.79 (1.35)	10.41 (0.92)	NS
Mesial to 5	3	8.75 (0.83)	7.56 (1.48)	S	8.61 (1.14)	8.00 (1.38)	NS
	5	9.21 (0.80)	8.60 (1.76)	S	9.10 (1.39)	9.05 (1.66)	NS
	7	9.18 (0.75)	9.02 (1.51)	NS	8.89 (1.56)	9.40 (2.03)	NS
Mesial to 4	3	8.97 (1.37)	8.49 (1.14)	NS	8.47 (1.06)	7.90 (1.36)	NS
	5	9.41 (1.23)	9.27 (1.48)	NS	8.83(0.91)	9.31(1.81)	NS
	7	8.98 (1.51)	9.24 (1.64)	NS	9.14 (1.57)	9.62 (2.26)	NS
Mesial to 3	3	7.48 (1.11)	6.92 (0.94)	NS	8.01(2.25)	7.35 (1.82)	NS
	5	9.19 (1.94)	7.93 (1.48)	NS	9.11(2.08)	8.32 (1.60)	NS
	7	9.47 (2.40)	7.52 (1.67)	S	9.07 (2.32)	8.07 (1.81)	NS
Mesial to 2	3	8.90 (2.50)	6.47 (0.90)	S	8.59 (1.93)	6.90 (2.88)	NS
	5	9.82(2.12)	7.10(0.65)	S	9.50(1.82)	6.75(1.74)	S
	7	10.12(1.97)	6.93(1.14)	S	9.72(1.99)	6.54(1.97)	S
Mesial to 1	3	7.49(1.68)	5.09(0.87)	S	7.49(1.68)	5.09(0.87)	S
	5	7.62(2.34)	5.89(1.40)	S	7.62(2.34)	5.89(1.40)	S
	7	7.90(2.68)	6.24(1.36)	NS	7.90(2.68)	6.24(1.36)	NS

R=Right, L=Left, Max=Maxillary, Mand=Mandibular, ABT= Alveolar Bone Thickness, SD=Standard Deviation, Males= 22, Sig.= significant at $P \leq 0.05$, S= significant, NS= non significant.

For females, ABT Max.- Mand. comparison [Table 4] displayed significant differences with the Max. ABT was higher than Mand. except in the following sites in which there were no significant differences: R and L Mesial to 7 at 5 and 7 mm levels; L Mesial to 6 at 7mm level and R and L Mesial to 5 at 7 mm levels.

DISCUSSION

Advanced CBCT systems; in the current study was NewTom CBCT machine; with high spatial resolution, sub-millimeter voxel sizes, small FOV, and a smaller focal spot are considered highly accurate in regard to linear measurements as stated by Sönmez *et al.* [28].

Table 4: Maxillary- mandibular comparison of alveolar bone thickness at different levels from cement-enamel junction for females

Variable	Level (mm)	R			L		
		Max. Mean (SD)	Mand. Mean (SD)	Sig.	Max. Mean (SD)	Mand. Mean (SD)	Sig.
Mesial to 7	3	13.70 (0.97)	11.40 (1.61)	S	13.70 (0.97)	11.86 (2.51)	S
	5	14.41(1.09)	13.22 (1.78)	NS	14.41(1.09)	13.73 (2.08)	NS
	7	14.17 (1.19)	12.13 (2.63)	NS	14.17 (1.19)	14.46 (1.62)	NS
Mesial to 6	3	10.75 (1.06)	9.00 (0.96)	S	10.84 (0.71)	8.67 (1.19)	S
	5	11.18 (0.70)	9.82 (1.02)	S	11.17 (0.73)	10.09 (1.31)	S
	7	11.43 (0.86)	10.63 (1.05)	S	10.98 (0.95)	10.90 (1.44)	NS
Mesial to 5	3	8.89 (1.30)	7.46 (0.92)	S	9.18 (0.98)	7.27 (0.81)	S
	5	9.59 (1.10)	8.22 (1.29)	S	9.86 (1.11)	8.21 (1.29)	S
	7	9.90 (0.99)	8.90 (1.46)	NS	9.96 (1.18)	8.93 (1.59)	NS
Mesial to 4	3	8.92 (1.26)	7.31(1.63)	S	8.88 (1.20)	7.16 (1.05)	S
	5	10.36 (1.18)	8.43 (1.62)	S	9.84 (1.19)	8.16 (1.18)	S
	7	11.33 (2.05)	8.74 (1.58)	S	10.06 (1.73)	8.55 (1.46)	S
Mesial to 3	3	8.24 (1.40)	6.84 (1.31)	S	8.89 (1.87)	6.62 (1.16)	S
	5	9.82 (2.24)	7.59 (1.38)	S	10.19 (2.07)	7.35 (1.18)	S
	7	10.57 (2.61)	7.56 (1.57)	S	10.64 (2.61)	7.42 (1.46)	S
Mesial to 2	3	8.76 (2.21)	6.26 (0.79)	S	9.18 (1.97)	6.03 (0.68)	S
	5	10.61 (2.77)	6.86 (0.67)	S	10.06 (2.28)	6.77 (1.13)	S
	7	11.16 (2.83)	6.87 (0.74)	S	10.49 (1.77)	6.55 (1.44)	S
Mesial to 1	3	7.69 (1.03)	5.46 (0.78)	S	7.69 (1.03)	5.46 (0.78)	S
	5	7.95 (1.09)	5.73 (1.01)	S	7.95 (1.09)	5.73 (1.01)	S
	7	8.27 (1.40)	6.35 (1.09)	S	8.27 (1.40)	6.35 (1.09)	S

R=Right, L=Left,Max=Maxillary, Mand=Mandibular, ABT= Alveolar Bone Thickness, SD=Standard Deviation, Females=26. Sig = significant at $P \leq 0.05$, S= significant , NS= non significant.

Safe MOPs with no trauma to the neighboring anatomical structures such as the roots, nasal cavity, Max. sinus, blood vessels or nerves are among the most important factors to be considered in the selection of MOP sites. Likewise, both the quality and quantity of the AB play a vital role in the success of MOP placement. Sangsuwon *et al.* [21] gave general guidelines to perform MOPs that can be performed in both the buccal and lingual cortical plates around the target tooth to encourage more bone remodeling with the buccal cortical plate is the most favorable site for placement of MOPs. Thus, in the current study different sites at different levels from both buccal and palatal/lingual sides were measured

to construct more detailed guidelines for the orthodontist when performing MOPs at different Max.-Mand. locations.

Three different levels from CEJ were selected to find the maximum ABT available to perform MOPs. According to Sangsuwon *et al.*, perforation depth of MOPs can be increased to compensate for the smaller number of perforations when a higher number of MOPs is not feasible. The number and depth of MOPs affect the inflammatory response and thus turnover of the bone as proved by Fokas *et al.*[29]. In the current study, CEJ was used for this purpose due to its constant position, easy access, and visibility by the examiner. Other studies measuring ABT depended on different anatomical landmarks as reference points such as the alveolar crest [30] or CEJ [31-33] as they used their corresponding software to measure ABT at variables levels apical to the CEJ.

The recommended penetration depths of MOPs should be in the range of 3–7 mm into the bone as suggested by Sangsuwon *et al.* [21]. In the current study, the lowest mean ABT; for proposed MOPs; was at Mand. anterior teeth at a 3mm level from CEJ, which was greater than the range recommended by Sangsuwon *et al.* [21].

The trend for the increase in ABT, i.e., increasing from the anterior toward the molar region was the same as in Katranji *et al.*[34] cadaver study in edentulous and dentate regions of elderly in different regions correlating to molar, premolar, and anterior teeth, however, Katranji *et al.* [34] determined buccal and lingual cortical bone plates thickness separately instead of total ABT as in the present study. A possible explanation for this trend may be attributed to heritability estimates of ABT, especially around Max. and Mand. incisors. According to

Goshtasbi *et al.* [35], genetic factors played significant roles in determining ABT around these teeth which is moderate to high effect.

The tendency toward increasing Max. and Mand. ABT gradually from inter-incisal distally toward molar and from near CEJ toward apical regions, came in agreement with the general findings of Golshah *et al.* [36], who found that optimal ABT varies depending on the sagittal skeletal pattern of Persian adults for miniscrew insertion and that the maximum ABT in the Max. and the Mand. in class I patients were posteriorly at the site of first and second molar at 2, 4 and 6 mm levels and were higher in Max. than Mand.

The general trend in the current study was increased ABT in Max. than Mand., this came in contrast to Khumsarn *et al.* [31] findings as they found; among the measured variables; in their comparative study of Class I and Class III sagittal skeletal patterns of Thai patients, Mand. alveolar process was thicker and wider in Class I patients than Max. process. Likewise, Coşkun and Kaya [33] results of ABT (the combined buccal cortical bone, cancellous bone, and lingual cortical bone thicknesses) disagreed with current findings in that ABT in the Mand. was higher than in Max. Likewise, there is disagreement with the findings of above mentioned Golshah *et al.* (2021) [36] study. Similarly, in Andrews *et al.* [27] study, ABT in untreated optimal occlusions was more in the Mand. than in Max. Such differences in the outcomes may be attributed to different sample populations, different settings and resolution of the CBCT system, different voxel sizes used for the CBCT measurements, different software accuracy used, or using different anatomical landmarks as reference points for determination of ABT and suspected subjective variations in landmark identification and variable measurement.

From the above mentioned results, the tested hypothesis was rejected as there were gender and Max.-Mand. differences in ABT except for specific locations.

CONCLUSIONS

Detailed information for ABT concerning MOP sites was obtained at different locations of both dental arches for both genders. There was a general trend for increasing Max. and Mand. ABT gradually from the inter-incisal region distally toward molars and from near CEJ toward apical regions. With abundant ABT was found apically at the Max. posterior region, while, the least ABT was found cervically at Mand. anterior region.

REFERENCES

- 1- Mitwally RA, Alesawi LM, Arishi TQ, Humedi AY, Al Baaltahin SS, Saeedi YA, Al Murdif AS, Al Zuhayr AZ, Albilali HW, Habis RM, Bakhsh MA. Factors affecting orthodontic treatment time and how to predict it. *Int J Community Med Public Health* 2022;9(1):492-6. <https://doi.org/10.18203/2394-6040.ijcmph20214798>.
- 2- Abbing A, Koretsi V, Eliades T, Papageorgiou SN. Duration of orthodontic treatment with fixed appliances in adolescents and adults : a systematic review with meta-analysis. *Prog Orthod* 2020; 2(1):1:37. <https://doi.org/10.1186/s40510-020-00334-4>
- 3- Bhad Patil WA, Karemore AA. Efficacy of pulsed electro-magnetic field in reducing treatment time: A clinical investigation. *Am J Orthod Dentofacial Orthop* 2022; 161(5):652-8. <https://doi.org/10.1016/j.ajodo.2020.12.025>

- 4- Jheon AH, Oberoi S, Solem RC, Kapila S. Moving towards precision orthodontics: An evolving paradigm shift in the planning and delivery of customized orthodontic therapy. *Orthod Craniofac Res* 2017;20 Suppl 1:106-13. <https://doi.org/10.1111/ocr.12171>
- 5- Jahanbin A, Hasanzadeh N, Khaki S, Shafae H. Comparison of self-ligating Damon3 and conventional MBT brackets regarding alignment efficiency and pain experience: A randomized clinical trial. *J Dent Res Dent Clin Dent Prospects* 2019; 13(4): 281-8. <https://doi.org/10.15171/joddd.2019.043>
- 6- Bhat SS, Sharma Y, Bagaria A, Sharma S, Marwaha J, Verma I. Drug induced orthodontic tooth movement: A comprehensive review. *SSRG-IJMS* 2019; 6(11):11 –4. <http://dx.doi.org/10.14445/23939117/IJMS-V6I11P104>
- 7- Bakdach WMM, Hadad R. Effectiveness of low-level laser therapy in accelerating the orthodontic tooth movement: A systematic review and meta-analysis. *Dent Med Probl* 2020; 57(1):73-94. <https://doi.org/10.17219/dmp/112446>
- 8- El-Bialy T, Farouk K, Carlyle TD, Wiltshire W, Drummond R, Dumore T, Knowlton K, Tompson B. Effect of low intensity pulsed ultrasound (LIPUS) on tooth movement and root resorption: A prospective multi-center randomized controlled trial. *J Clin Med* 2020; 9:804. <https://doi.org/10.3390/jcm9030804>
- 9- Li Y, Deng S, Mei L, Li J, Qi M, Su S, Li Y, Zheng W. Accuracy of alveolar bone height and thickness measurements in cone beam computed tomography:a systematic review and meta-analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2019; 128(6):667-79.

10- Gil APS, Haas OL Jr and M´endez-Manj´on I, Masiá-Gridilla J, Valls-Ontañón A, Hernández-Alfaro F , Guijarro-Martínez R. Alveolar corticotomies for accelerated orthodontics: a systematic review. *J Craniomaxillofac Surg* 2018;46(3):438–45. <https://doi.org/10.1016/j.jcms.2017.12.030>

11- Ji Y, Tang Y , Wu Q , Huang D , Zhu J, Kang F. The effects of mandibular osteotomy on maxillary orthodontic tooth movement and bone remodelling in a rat model. *Eur J Orthod* 2021;43(4): 467–72. <https://doi.org/10.1093/ejo/cjaa053>

12- Liu R, Huang L, Xiao X, Guan Y, Jiang Y, Yin X, Zou S, Ye Q. Biomechanical interfaces of corticotomies on periodontal tissue remodeling during orthodontic tooth movement. *Coatings* 2021; 11(1):1. <https://doi.org/10.3390/coatings11010001>

13- Figueiredo D, Houara R, Pinto L, Diniz A, de Araújo V, Thabane L, Soares R, Oliveira D. Effects of piezocision in orthodontic tooth movement: A systematic review of comparative studies. *J Clin Exp Dent* 2019;11(11):e1078-92. <https://doi.org/10.4317/jced.56328>

14- Alikhani M, Raptis M, Zoldan B, Sangsuwon C, Lee YB, Alyami B, Corpodian C, Barrera LM, Alansari S, Khoo E, Teixeira C. Effect of micro-osteoperforations on the rate of tooth movement. *Am J Orthod Dentofacial Orthop* 2013;14(5)4:639-48. <https://doi.org/10.1016/j.ajodo.2013.06.017>

15- Hoffmann S, Papadopoulos N, Visel D, Visel T, Jost-Brinkmann PG, Präger TM. Influence of piezotomy and osteoperforation of the alveolar process on the rate of orthodontic tooth movement: a systematic review. *J Orofac Orthop* 2017;78(4):301–11. <https://doi.org/10.1007/s00056-017-0085-1>

16- Isola G, Matarese G, Cordasco G, Perillo L and Ramaglia L. Mechanobiology of the tooth movement during the orthodontic treatment: a literature review. *Minerva Stomatologica* 2016; 65(5):299–327.

17- Alansari S, Sangsuwon C, Vongthongleur T, Kwal R, Teo M, Lee YB, Nervina J, Teixeira C, Alikhani M. Biological principles behind accelerated tooth movement. *Semin Orthod* 2015; 21, 151–61. <http://dx.doi.org/10.1053/j.sodo.2015.06.001>

18- Charavet C, Lambert F, Lecloux G, Le Gall M. Accelerated orthodontic treatment using corticotomies: what are the minimally invasive alternatives? *Orthod Fr* 2019;90(1):5–12.

19- Alikhani M, Alansari S, Sangsuwon C, Alikhani M, Chou MY, Alyami B, Nervina JM, Teixeira CC. Micro-osteoperforations: Minimally invasive accelerated tooth movement. *Semin Orthod* 2015; 21:162–9. <http://dx.doi.org/10.1053/j.sodo.2015.06.002>

20- Maspero C, Cappella A, Dolci C, Cagetti MG, Inchingolo F, Sforza C. Is orthodontic treatment with microperforations worth it? A scoping review. *Children (Basel)* 2022; 9(2): 208. <https://doi.org/10.3390/children9020208>

21- Sangsuwon C, Alansari S, Nervina J, Teixeira CC, Alikhani M. Micro-osteoperforations in accelerated orthodontics. *Clin Dent Rev* 2018; 2:4. <https://doi.org/10.1007/s41894-017-0013-1>

22- Aboalnaga AA, Fayed MMS, El-Ashmawi NA, Soliman SA. Effect of micro-osteoperforation on the rate of canine retraction: a split-mouth randomized controlled trial. *Prog Orthod* 2019; 3;20(1):21. <https://doi.org/10.1186/s40510-019-0274-0>

- 23- Alqadasi B, Aldhorae K, Halboub E, Mahgoub N, Alnasri A, Assiry A, Xia HY. The Effectiveness of micro-osteoperforations during canine retraction: A three-dimensional randomized clinical trial. *J Int Soc Prev Community Dent* 2019; 9(6):637-45. https://doi.org/10.4103/jispcd.jispcd_233_19
- 24- Fattoria L, Sendyka M, de Paivab JB; Normandoc D, Netod JR. Micro-osteoperforation effectiveness on tooth movement rate and impact on oral health related quality of life: A randomized clinical trial. *Angle Orthod* 2020;90(5):640–7. <https://doi.org/10.2319/110819-707.1>
- 25- Mittal R, Attri S, Batra P, Sonar S, Sharma K, Raghavan S. Comparison of orthodontic space closure using micro-osteoperforation and passive self-ligating appliances or conventional fixed appliances: A randomized controlled trial. *Angle Orthod.* 2020; 90(5):634-9. <https://doi.org/10.2319/111119-712.1>
- 26- Shahabee M, Shafae H, Abtahi M, Rangrazi A, Bardideh E. Effect of micro-osteoperforation on the rate of orthodontic tooth movement—a systematic review and a meta-analysis. *Eur J Orthod* 2020; 42(2):211-21. <https://doi.org/10.1093/ejo/cjz049>
- 27- Andrews WA, Abdulrazzaqb WS, Hunt JE, Mendesd LM, Hallmane LA. Incisor position and alveolar bone thickness: A comparative analysis of two untreated samples using lateral cephalograms. *Angle Orthod* 2022;92(1):3–10. <https://doi.org/10.2319/022320-122.1>
- 28- Sönmez G, Koç C, Kamburoğlu K. Accuracy of linear and volumetric measurements of artificial ERR cavities by using CBCT

images obtained at 4 different voxel sizes and measured by using 4 different software:an ex vivo research. *Dentomaxillofac Radiol* 2018;47(8):20170325. <https://doi.org/10.1259/dmfr.20170325>

29- Fokas G, Vaughn VM, Scarfe WC, Bornstein MM. Accuracy of linear measurements on CBCT images related to presurgical implant treatment planning: a systematic review. *Clin Oral Implants Res* 2018;29 Suppl (16):393-415. <https://doi.org/10.1111/clr.13142>

30- Sangswon C, Alansari S, Lee Y, Nervina J, Alikhani M. Step-by-step guide for performing micro-osteoperforations. Chapter 6:99-116. Springer international publishing AG 2017. M Alikhani(ed.), *Clinical Guide to Accelerated Orthodontics*. <https://doi.org/10.1007/s41894-017-0013-1>

31- Khumsarn N, Patanaporn V, Janhom A, Jotikasthira D. Comparison of interradicular distances and cortical bone thickness in Thai patients with Class I and Class II skeletal patterns using cone-beam computed tomography. *Imaging Sci Dent* 2016;46(2):117–25. <https://doi.org/10.5624/isd.2016.46.2.117>

32- Shrestha B, Kurtzman GM, Basnet BB, Adhikari G, Rokaya D. Assessment of buccal bone thickness of aesthetic maxillary region in Nepalese adults: A Cone Beam Computed Tomography (CBCT) Study. *JNPS* 2019; 2(2): 63-72. <http://dx.doi.org/10.3126/jnprosoc.v2i2.31192>

33- Coşkun İ, Kaya B. Relationship between alveolar bone thickness, tooth root morphology, and sagittal skeletal pattern: A cone beam computed tomography study. *J Orofac Orthop* 2019; 80(3):144-58. <https://doi.org/10.1051/orthodfr/2019002>

- 34- Katranji A, Misch K, Wang H-L. Cortical bone thickness in dentate and edentulous human cadavers. *J Periodontol* 2007; 78 (5): 874-8. <https://doi.org/10.1902/jop.2007.060342>
- 35- Goshtasbi N, Hazzard J, Mehrkhodavandi N, Al-Qawasmia R. The heritability of alveolar bone thickness in siblings seeking orthodontic treatment. *Am J Orthod Dentofacial Orthop* 2020;157(6):803-8. <https://doi.org/10.1016/j.ajodo.2019.06.019>
- 36- Golshah A, Salahshour M, Nikkerdar N. Interradicular distance and alveolar bone thickness for mini-screw insertion: a CBCT study of Persian adults with different sagittal skeletal patterns. *BMC Oral Health* 2021; 21(1):534. <https://doi.org/10.1186/s12903-021-01891-8>